

ISOLATION AND STRUCTURE ELUCIDATION
OF NATURAL PRODUCTS FROM MARINE
ORGANISMS AND DETERMINATION OF
THEIR BIOLOGICAL ACTIVITY

A THESIS SUBMITTED IN PARTIAL PULFILMENT OF THE
REQUIREMENTS FOR THE DEGREE

OF

MASTER OF PHILOSOPHY

OF THE


UNIVERSITY OF COLOMBO

SRI LANKA

BY

447268

JAYAKODY ARACHCHIGE CHANDRASIRI



DEPARTMENT OF CHEMISTRY
UNIVERSITY OF COLOMBO
COLOMBO 03

JULY 1990


ABSTRACT




Marine organisms are a fascinating source of biologically active secondary metabolites such as anti-bacterial, anti-fungal, insecticidal, cytotoxic and anti-cancer agents.

This research project was aimed at identifying such bio-active compounds from Sri Lankan marine organisms. The extracts of a number of organisms collected from the Sri Lankan coastal waters were screened for anti-bacterial (against Staphylococcus aureus, E. coli, and Pseudomonas sp.) and anti-fungal (against Cladosporium cladosporioides) activity.

Chemical investigation of the MeOH-CH₂Cl₂ (1:1) extract of the soft coral Sinularia abrupta collected at Kalkudah afforded two new closely related furano-cembranoid diterpenes (246) and (247). The structures of these compounds were deduced on the basis of their spectral data.




Chromatographic separation of the MeOH-CH₂Cl₂ (1 : 1) extract of the soft coral Sinularia crispera collected at Unawatuna yielded 1-O-octadecyl glycerol (batyl alcohol) (248) which showed potent anti-bacterial activity against Pseudomonas sp.




(248)

A specimen of the soft coral Sinularia abrupta collected from Unawatuna, afforded two bio-active constituents ; a diacyl glyceryl ether (249) and the furano-sesquiterpene, methyl (5', E) -5-(2', 6'-dimethylocta-5', 7'-dienyl) furan - 3 - carboxylate (44).




(249)



(44)

Activity directed fractionation of the CHCl₃ extract of an unidentified sponge collected at Wellawatta

furnished two furano-sesterpenes, which showed strong anti-bacterial activity against Staphylococcus aureus. One of them was identified as variabilin (228).



(228)