

**158** Katulanda, P; Shine, B; Katulanda, GW; Mahesh, JG; Silva, A; Ganegoda, UA; **Sheriff, MHR**; Somasundaram, N; Wijesuriya, M; Matthews, DR

Prevalence and phenotype characteristics of latent autoimmune diabetes in adulthood (LADA) and type 1 diabetes among young adults in Sri Lanka; Abstract; Sri Lanka Medical Association - 121st Annual Scientific Sessions ; 2008\_.46pp

Abstract : Background: Sri Lankans are at high risk of diabetes. Knowledge of the aetiology is useful in disease prevention. Objectives: To determine the prevalence and clinico-metabolic characteristics of Glutamic Acid Decarboxylase Antibody (GAD) positive diabetes among young adult-onset subjects in Sri Lanka. Design, setting and methods: We recruited a sample of 1007 consecutive diabetic subjects 45 years (age of diagnosis 16 to 40 years). Clinical and metabolic characterization was performed. GAD antibodies were measured using a radioimmunoassay method. A titre  $>14\text{U/L}$  (WHO 97.5 percentile) was considered positive. LADA was diagnosed when subjects were GAD+ve,  $>30$  years and insulin independent  $>6$  months from diagnosis. Insulin resistance assessed using homeostatic model (HOMA-IR). Data were analysed using SPSS. Results: Mean age of the cohort was 36.7 years ( $\pm 5.8$ ), males 42.3%. Overall 16% were insulin treated. GAD was positive in 5.4% ( $n=54$ ) of our sample. The GAD titre had a significant negative correlation with the time of commencement of insulin ( $r=-0.22$ ,  $p=0.006$ ). The GAD+ve type 1 diabetes (T1DM) and LADA were 2.1% and 2.6% respectively. Among subjects with T1DM, LADA and type 2 diabetes (T2DM), BMI ( $\text{kg/m}^2$ ) [19.6, 22.2 and 25.0], fasting C-peptide ( $\text{nmol/l}$ ) [0.08, 0.52 and 0.65] and HOMAIR [0.28, 2.0 and 1.75] showed significant differences ( $p<0.001$ ). Metabolic syndrome was present in 0%, 46% and 67.3% of the T1DM, LADA and T2DM subjects respectively ( $p<0.001$ ). Conclusions: GAD positive diabetes is less prevalent among young adult diabetic subjects in Sri Lanka compared to T2DM. Subjects with LADA are phenotypically more related to T2DM than to T1DM.